

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
25 October 2001 (25.10.2001)

PCT

(10) International Publication Number
WO 01/78741 A1

(51) International Patent Classification⁷: **A61K 31/575**,
31/46, A61P 11/06 // (A61K 31/575, 31:46)

(21) International Application Number: PCT/GB01/01641

(22) International Filing Date: 11 April 2001 (11.04.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
0009583.6 18 April 2000 (18.04.2000) GB

(71) Applicant (for all designated States except US): **GLAXO GROUP LIMITED** [GB/GB]; Glaxo Wellcome House, Berkeley Avenue, Greenford, Middlesex UB6 0NN (GB).

(72) Inventor; and

(75) Inventor/Applicant (for US only): **GAVIN, Brian, Charles** [IE/IE]; GlaxoSmithKline, P.O. Box 700, Grange Road, Rathfarnham, 16 Dublin (IE).

(74) Agent: **LEAROYD, Stephanie, Anne**; GlaxoSmithKline, Corporate Intellectual Property, Two New Horizons Court, Brentford, Middlesex TW8 9EP (GB).

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

- with international search report
- before the expiration of the time limit for amending the claims and to be republished in the event of receipt of amendments

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: **MEDICAL COMBINATIONS COMPRISING TIOTROPIUM AND BUDESONIDE**

(57) Abstract: The present invention is concerned with pharmaceutical formulations comprising a combination of tiotropium and budesonide and the use of such formulations in medicine, particularly in the prophylaxis and treatment of respiratory diseases.

WO 01/78741 A1

MEDICAL COMBINATIONS COMPRISING TIOTROPIUM AND BUDESONIDE

The present invention is concerned with combinations of tiotropium and budesonide, particularly compositions containing a combination of tiotropium and budesonide and the use of such compositions in medicine, particularly in the prophylaxis and treatment of respiratory diseases.

Tiotropium i.e. $(1\alpha, 2\beta, 4\beta, 5\alpha, 7\beta)$ -7-[(hydroxydi-2-thienylacetyl)oxy]-9,9-dimethyl-3-oxa-9-azoniatricyclo[3.3.2.0]nonane and particularly its bromide salt is a well-known anti-cholinergic agent, described in EP418,716 for the treatment of bronchial asthma and related disorders.

DE 2,323,215 and US 3,929,768 describe budesonide i.e. $(11\beta, 16\alpha)$ -16,17-[butylidenebis(oxy)]-11,21-dihydroxypregna-1,4-diene-3,20-dione, salts thereof and pharmaceutical formulations thereof. Budesonide is an antiinflammatory corticosteroid, which is now used clinically in the treatment of bronchial asthma and related disorders.

Although tiotropium bromide and budesonide are effective therapies, there exists a clinical need for asthma therapies having potent and selective action and having an advantageous profile of action.

Therefore, according to the present invention there is provided a combination of tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof.

It will be appreciated that the compounds of the combination may be administered simultaneously, either in the same or different pharmaceutical formulations or sequentially. If there is sequential administration, the delay in administering the second compound should not be such as to lose the beneficial therapeutic effect of the combination.

According to a further aspect of the present invention, there is provided a pharmaceutical formulation comprising tiotropium or a pharmaceutically

acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients. According to
5 a preferred aspect of the present invention, there is provided a pharmaceutical formulation comprising tiotropium bromide and budesonide, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients. In the most preferred aspect, the above pharmaceutical formulations are suitable for administration by inhalation.

10 It is to be understood that the present invention covers all combinations of particular and preferred aspects of the invention described herein.

As would be appreciated by the skilled person, budesonide is sold as an epimer
15 mixture of the α - and β -propyl forms. The present invention includes each epimer of budesonide either in substantially pure form or admixed in any proportions.

By the term "physiologically functional derivative" is meant a chemical derivative
20 of tiotropium or budesonide having the same physiological function as the free compound, for example, by being convertible in the body thereto. According to the present invention, examples of physiologically functional derivatives include esters.

25 Suitable salts according to the invention include those formed with both organic and inorganic acids. Pharmaceutically acceptable acid addition salts include but are not limited to those formed from hydrochloric, hydrobromic, sulphuric, citric, tartaric, phosphoric, lactic, pyruvic, acetic, trifluoroacetic, succinic, oxalic, fumaric, maleic, oxaloacetic, methanesulphonic, ethanesulphonic, p-
30 toluenesulphonic, benzenesulphonic, isethionic, and naphthalenecarboxylic, such as 1-hydroxy-2-naphthalenecarboxylic acids.

Pharmaceutically acceptable esters of tiotropium or budesonide may have a
35 hydroxyl group converted to a C₁₋₆alkyl, aryl, aryl C₁₋₆ alkyl, or amino acid ester.

As mentioned above, both tiotropium and budesonide and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have been described for use in the treatment of respiratory diseases. Therefore, formulations of tiotropium and budesonide and their pharmaceutically acceptable salts, solvates, and physiologically functional derivatives have use in the prophylaxis and treatment of clinical conditions for which an anticholinergic agent and/or an antiinflammatory corticosteroid is indicated. Such conditions include diseases associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary diseases (COPD) (e.g. chronic and wheezy bronchitis, emphysema), respiratory tract infection and upper respiratory tract disease.

Accordingly, the present invention provides a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which an anticholinergic agent and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a combination of tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof. The present invention further provides a method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which an anticholinergic agent and/or antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a pharmaceutical formulation comprising tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient. In a preferred aspect, there is provided such a method which comprises administration of a therapeutically effective amount of a pharmaceutical formulation comprising tiotropium bromide and budesonide, and a pharmaceutically acceptable carrier or excipient. In particular, the present invention provides such methods for the prophylaxis or treatment of a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

In the alternative, there is provided a combination of tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, for use in therapy, particularly for use in the prophylaxis or treatment of a clinical condition for which an anticholinergic agent and/or antiinflammatory corticosteroid is indicated. In particular, there is provided a pharmaceutical formulation comprising tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof (suitably, tiotropium bromide) and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient for use in therapy, particularly for use in the prophylaxis or treatment of a clinical condition for which an anticholinergic agent and/or antiinflammatory corticosteroid is indicated. In a preferred aspect, the invention is concerned with the prophylaxis or treatment of a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

The amount of tiotropium and budesonide, or a pharmaceutically acceptable salt, solvate or physiologically functional derivative thereof which is required to achieve a therapeutic effect will, of course, vary with the particular compound, the route of administration, the subject under treatment, and the particular disorder or disease being treated. As a monotherapy, tiotropium bromide is generally administered to adult humans by aerosol inhalation at a dose of 10mcg to 200mcg once or twice daily. As a monotherapy, budesonide is generally administered to adult humans by aerosol inhalation at a dose of from 200mcg to 1.6mg daily, taken as 2 divided doses.

While it is possible for the active ingredients of the combination to be administered as the raw chemical, it is preferable to present them as a pharmaceutical formulation. When the individual compounds of the combination are administered separately, they are generally each presented as a pharmaceutical formulation as described previously in the art.

Pharmaceutical formulations are often prescribed to the patient in "patient packs" containing the whole course of treatment in a single package. Patient packs have an advantage over traditional prescriptions, where a pharmacist divides a patient's supply of a pharmaceutical from a bulk supply, in that the patient always has access to the package insert contained in the patient pack, normally missing in traditional prescriptions. The inclusion of a package insert has been shown to improve patient compliance with the physician's instructions and, therefore, lead generally to more successful treatment. It will be understood that the administration of the combination of the invention by means of a single patient pack, or patient packs of each component compound, and containing a package insert instructing the patient to the correct use of the invention is a desirable additional feature of the invention.

Hereinafter, the term "active ingredients" means tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, preferably tiotropium bromide, and budesonide, or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof.

Suitably, the pharmaceutical formulations which are suitable for inhalation according to the invention comprise the active ingredients in amounts such that each actuation provides therapeutically effective dose, for example, a dose of tiotropium of 10mcg to 200mcg, preferably 20mcg to 100mcg and a dose of budesonide of 100mcg to 1.6mg, preferably 200mcg to 1mg, more preferably, 200mcg to 400mcg.

The pharmaceutical formulations according to the invention may further include other therapeutic agents for example anti-inflammatory agents such as other corticosteroids (e.g. fluticasone propionate, beclomethasone dipropionate, mometasone furoate, or triamcinolone acetonide) or NSAIDs (e.g. sodium cromoglycate, nedocromil sodium, PDE-4 inhibitors, leukotriene antagonists, iNOS inhibitors, tryptase and elastase inhibitors, beta-2 integrin antagonists and adenosine 2a agonists), or β_2 -adrenoreceptor agonists (such as salbutamol, salmeterol, formoterol, fenoterol or terbutaline and salts thereof), or other anticholinergic agents (such as ipratropium).

The formulations include those suitable for oral, parenteral (including subcutaneous, intradermal, intramuscular, intravenous and intraarticular), intranasal, inhalation (including fine particle dusts or mists which may be generated by means of various types of metered dose pressurised aerosols, nebulisers or insufflators), rectal and topical (including dermal, buccal, sublingual and intraocular) administration although the most suitable route may depend upon for example the condition and disorder of the recipient. The formulations may conveniently be presented in unit dosage form and may be prepared by any of the methods well known in the art of pharmacy. All methods include the step of bringing the active ingredients into association with the carrier which constitutes one or more accessory ingredients. In general the formulations are prepared by uniformly and intimately bringing into association the active ingredients with liquid carriers or finely divided solid carriers or both and then, if necessary, shaping the product into the desired formulation.

Formulations for inhalation include powder compositions which will preferably contain lactose, and spray compositions which may be formulated, for example, as aqueous solutions or suspensions or as aerosols delivered from pressurised packs, with the use of a suitable propellant, e.g. dichlorodifluoromethane, trichlorofluoromethane, dichlorotetrafluoroethane, 1,1,1,2,3,3,3-heptafluoropropane, 1,1,1,2-tetrafluoroethane, carbon dioxide or other suitable gas. Suitable aerosol formulations include those described in EP 0372777 and WO93/11743. For suspension aerosols, the active ingredients should be micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the aerosol formulation, thus the active ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns, for example, 1 to 5 microns.

Intranasal sprays may be formulated with aqueous or non-aqueous vehicles with the addition of agents such as thickening agents, buffer salts or acid or alkali to adjust the pH, isotonicity adjusting agents or anti-oxidants.

Capsules and cartridges or for example gelatin, or blisters of for example laminated aluminium foil, for use in an inhaler or insufflator may be formulated

containing a powder mix of the active ingredients and a suitable powder base such as lactose or starch. In this aspect, the active ingredients are suitably micronised so as to permit inhalation of substantially all of the active ingredients into the lungs upon administration of the dry powder formulation, thus the active ingredients will have a particle size of less than 100 microns, desirably less than 20 microns, and preferably in the range 1 to 10 microns.

Solutions for inhalation by nebulation may be formulated with an aqueous vehicle with the addition of agents such as acid or alkali, buffer salts, isotonicity adjusting agents or antimicrobials. They may be sterilised by filtration or heating in an autoclave, or presented as a non-sterile product.

Preferred unit dosage formulations are those containing a pharmaceutically effective dose, as hereinbefore recited, or an appropriate fraction thereof, of the active ingredient. Thus, in the case of formulations designed for delivery by metered dose pressurised aerosols, one actuation of the aerosol may deliver half of the therapeutically effective amount such that two actuations are necessary to deliver the therapeutically effective dose.

It should be understood that in addition to the ingredients particularly mentioned above, the formulations of this invention may include other agents conventional in the art having regard to the type of formulation in question. Furthermore, the claimed formulations include bioequivalents as defined by the US Food and Drugs Agency.

For a better understanding of the invention, the following Examples are given by way of illustration.

EXAMPLESA: Metered Dose Inhalers

5 Example 1

	Per actuation
Tiotropium Bromide	100 microgram
Budesonide	200 microgram
1,1,1,2-Tetrafluoroethane	to 75.0mg

10 The micronised active ingredients are weighed into an aluminium can, 1,1,1,2-tetrafluoroethane is then added from a vacuum flask and a metering valve is crimped into place.

Similar methods may be used for the formulation of Examples 2 to 4:

15 Example 2

	Per actuation
Tiotropium Bromide	200 microgram
Budesonide	100 microgram
1,1,1,2-Tetrafluoroethane	to 75.0mg

Example 3

	Per actuation
Tiotropium Bromide	9 microgram
Budesonide	100 microgram
1,1,1,2-Tetrafluoroethane	to 75.0mg

Example 4

	Per actuation
Tiotropium Bromide	18 microgram
Budesonide	100 microgram
1,1,1,2-Tetrafluoroethane	to 75.0mg

B: Dry Powder Inhalers

5

Example 5

	Per cartridge or blister
Tiotropium Bromide	100 microgram
Budesonide	200 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

10

The active ingredients are micronised and bulk blended with the lactose in the proportions given above. The blend is filled into hard gelatin capsules or cartridges or in specifically constructed double foil blister packs to be administered by an inhaler such as a Rotahaler, Diskhaler, or Diskus inhaler (each of these being a Trademark of Glaxo Group Limited).

15

Similar methods may be used for the formulations of Example 6 to 8:

Example 6

	Per cartridge or blister
Tiotropium Bromide	200 microgram
Budesonide	100 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

20

Example 7

	Per cartridge or blister
Tiotropium Bromide	9 microgram
Budesonide	100 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

Example 8

5

	Per cartridge or blister
Tiotropium Bromide	18 microgram
Budesonide	100 microgram
Lactose Ph. Eur.	to 12.5mg or to 25.0mg

Claims

1. A pharmaceutical formulation comprising tiotropium or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof and budesonide or a pharmaceutically acceptable salt, solvate, or physiologically functional derivative thereof, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.
2. A pharmaceutical formulation comprising tiotropium bromide and budesonide, and a pharmaceutically acceptable carrier or excipient, and optionally one or more other therapeutic ingredients.
3. A pharmaceutical formulation according to claim 1 or 2 which is suitable for administration by inhalation.
4. A pharmaceutical formulation according to any of claims 1 to 3 wherein the pharmaceutically acceptable carrier or excipient is lactose.
5. A pharmaceutical formulation according to any of claims 1 to 3 wherein the pharmaceutically acceptable carrier or excipient comprises 1,1,1,2-tetrafluoroethane and/or 1,1,1,2,3,3,3- heptafluoropropane.
6. A method for the prophylaxis or treatment of a clinical condition in a mammal, such as a human, for which an anticholinergic agent and/or an antiinflammatory corticosteroid is indicated, which comprises administration of a therapeutically effective amount of a pharmaceutical formulation according to any one of claims 1 to 5.
7. A method according to claim 6 wherein the clinical condition is a disease associated with reversible airways obstruction such as asthma, chronic obstructive pulmonary disease (COPD), respiratory tract infection or upper respiratory tract disease.

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/01641

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K31/575 A61K31/46 A61P11/06 //(A61K31/575,31:46)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, MEDLINE, BIOSIS, CHEM ABS Data, EMBASE, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 418 716 A (BOEHRINGER INGELHEIM KG ;BOEHRINGER INGELHEIM INT (DE)) 27 March 1991 (1991-03-27) cited in the application claims ---	1-7
Y	QURESHI F ET AL: "Effect of nebulized ipratropium on the hospitalization rates of children with asthma." NEW ENGLAND JOURNAL OF MEDICINE, (1998 OCT 8) 339 (15) 1030-5. , XP001007631 abstract --- -/--	1-7

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
- *Z* document member of the same patent family

Date of the actual completion of the international search

9 August 2001

Date of mailing of the international search report

04/09/2001

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel (+31-70) 340-2040, Tx. 31 651 epo nl,

Authorized officer

INTERNATIONAL SEARCH REPORT

Inte 1st Application No

PCT7GB 01/01641

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	BACULARD A: "Bronchodual in the long-term treatment of children with asthma." ARCHIVES DE PEDIATRIE, vol. 2, no. SUPPL. 2, 1995, pages 149S-153S, XP000914115 ISSN: 0929-693X abstract	1-7
Y	DE 23 23 215 A (BOFORS AB) 29 November 1973 (1973-11-29) cited in the application claims 1-24	1-7
Y	US 3 929 768 A (BRATTSAND RALPH LENNART ET AL) 30 December 1975 (1975-12-30) cited in the application claims 1-29	1-7
Y	BOWLER S: "LONG ACTING BETA AGONISTS" AUSTRALIAN FAMILY PHYSICIAN, XX, XX, vol. 27, no. 12, December 1998 (1998-12), pages 1115,1117-1118, XP000973076 the whole document	1-7
Y	BARNES P J ET AL: "EFFICACY OF INHALED CORTICOSTEROIDS IN ASTHMA" JOURNAL OF ALLERGY AND CLINICAL IMMUNOLOGY, MOSBY - YEARLY BOOK, INC, US, vol. 102, no. 4, 1998, pages 531-538, XP000913470 ISSN: 0091-6749 page 536, right-hand column, line 1-6 abstract	1-7
Y	O'CONNOR B J: "COMBINATION THERAPY" PULMONARY PHARMACOLOGY AND THERAPEUTICS, ACADEMIC PRESS, NEW YORK, NY, US, vol. 11, no. 5/6, 1998, pages 397-399, XP000911059 ISSN: 1094-5539 the whole document	1-7

INTERNATIONAL SEARCH REPORT

Intel Application No

PCT/GB 01/01641

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
EP 0418716	A	27-03-1991	DE 3931041 A	28-03-1991
			AT 103914 T	15-04-1994
			AU 642913 B	04-11-1993
			AU 6431890 A	18-04-1991
			BG 61295 B	30-04-1997
			CA 2066248 A,C	17-03-1991
			CZ 9004523 A	11-11-1998
			DD 297647 A	16-01-1992
			DE 59005250 D	11-05-1994
			DK 418716 T	02-05-1994
			WO 9104252 A	04-04-1991
			ES 2052125 T	01-07-1994
			HR 940723 A	30-06-1997
			HU 60740 A	28-10-1992
			HU 208823 B	28-01-1994
			HU 210612 B	29-05-1995
			IE 903342 A	10-04-1991
			IL 95691 A	23-07-1996
			JP 7030074 B	05-04-1995
			JP 5502438 T	28-04-1993
			KR 168432 B	15-01-1999
			MX 9203150 A	01-07-1992
			NO 301478 B	03-11-1997
			NZ 235306 A	24-06-1997
			PL 168468 B	29-02-1996
			PT 95312 A,B	22-05-1991
			SI 9011744 A,B	31-10-1997
			SK 452390 A	04-11-1998
			RU 2073677 C	20-02-1997
			US 5610163 A	11-03-1997
			ZA 9007338 A	26-08-1992
DE 2323215	A	29-11-1973	SE 378109 B	18-08-1975
			AT 328630 B	25-03-1976
			AT 436573 A	15-06-1975
			AU 5525373 A	07-11-1974
			BE 799727 A	17-09-1973
			CA 1002938 A	04-01-1977
			CH 595400 A	15-02-1978
			CS 178129 B	31-08-1977
			CY 1013 A	23-11-1979
			DD 104295 A	05-03-1974
			DK 134783 B	17-01-1977
			ES 414673 A	01-07-1976
			FI 50631 B	02-02-1976
			FR 2185405 A	04-01-1974
			GB 1429922 A	31-03-1976
			HK 49179 A	27-07-1979
			HU 166680 B	28-05-1975
			IL 42155 A	30-06-1977
			JP 1033476 C	20-02-1981
			JP 49041378 A	18-04-1974
			JP 55021760 B	12-06-1980
			KE 2970 A	20-07-1979
			NL 7306978 A,B,	21-11-1973
			NO 139640 B	08-01-1979
			SU 470954 A	15-05-1975

INTERNATIONAL SEARCH REPORT

International Application No

PCT/GB 01/01641

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
DE 2323215 A		US 3983233 A	28-09-1976
		YU 129173 A,B	28-02-1981
		ZA 7302955 A	24-04-1974
US 3929768 A	30-12-1975	SE 378109 B	18-08-1975
		AT 328630 B	25-03-1976
		AT 436573 A	15-06-1975
		AU 5525373 A	07-11-1974
		BE 799727 A	17-09-1973
		CA 1002938 A	04-01-1977
		CH 595400 A	15-02-1978
		CS 178129 B	31-08-1977
		CY 1013 A	23-11-1979
		DD 104295 A	05-03-1974
		DE 2323215 A	29-11-1973
		DK 134783 B	17-01-1977
		ES 414673 A	01-07-1976
		FI 50631 B	02-02-1976
		FR 2185405 A	04-01-1974
		GB 1429922 A	31-03-1976
		HK 49179 A	27-07-1979
		HU 166680 B	28-05-1975
		IL 42155 A	30-06-1977
		JP 1033476 C	20-02-1981
		JP 49041378 A	18-04-1974
		JP 55021760 B	12-06-1980
		KE 2970 A	20-07-1979
		NL 7306978 A,B,	21-11-1973
		NO 139640 B	08-01-1979
		SU 470954 A	15-05-1975
		US 3983233 A	28-09-1976
		YU 129173 A,B	28-02-1981
		ZA 7302955 A	24-04-1974